

Erratum

Erratum to “Folding degrees of azurins and pseudoazurins.  
Implications for structure and function”  
[Comput. Biol. Chem. 29 (2005) 345–353]

Ernesto Estrada<sup>a,\*</sup>, Eugenio Uriarte<sup>b</sup>

<sup>a</sup> Complex Systems Research group, X-ray Unit, Edificio CACTUS, University of Santiago de Compostela, 15782 Santiago de Compostela, Spain

<sup>b</sup> Department of Organic Chemistry, Faculty of Pharmacy University of Santiago de Compostela, 15782 Santiago de Compostela, Spain

Received 16 May 2007; accepted 16 May 2007


On page 346, the **T** matrix is defined as a diagonal matrix with diagonal entries equal to  $T_{ii} = \cos \theta_i$ , where  $\theta_i$  are the corresponding dihedral (torsion) angles of the protein backbone. However, in this work we have used in all calculations the function

$$f_i = \frac{1}{2}[1 + \text{sgn}(\cos \theta_i)] \cos \theta_i \quad (1)$$

where  $\text{sgn}(x)$  is the sign function defined as

$$\text{sgn}(x) = \begin{cases} -1, & x < 0 \\ 0, & x = 0 \\ 1, & x > 0 \end{cases} \quad (2)$$

The plot of this function is given below for illustration.



DOI of original article: [10.1016/j.combiolchem.2005.06.005](https://doi.org/10.1016/j.combiolchem.2005.06.005).

\* Corresponding author. Tel.: +34 981563100; fax: +34 981547077.

E-mail address: [estrada66@yahoo.com](mailto:estrada66@yahoo.com) (E. Estrada).